


Marcos A. G. García





# About me



PhD, inflation, susy

Minneapolis


Houston

Mexico City



RICE

Postdoc, CMB, GWs




PhD, math. phys.



Postdoc, dark matter

Madrid






Me a few hours ago...




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Solid evidence for Dark Matter







# 1. Beyond WIMPs



# 2. Inflation & reheating




# 3. FIMPs



# 4. Compact objects



# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating





# 3. FIMPs



# 4. Compact objects



# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating






# 3. FIMPs



# 4. Compact objects



# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating







# 3. FIMPs



# 4. Compact objects



# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating







# 3. FIMPs



# 4. Compact objects



# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating










# 3. FIMPs



# 4. Compact objects



# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating




# 3. FIMPs



# 4. Compact objects



# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects



# 5. Prospects

## The many virtues of the WIMP



$$\Omega_{\text{DM}} h^2 \equiv \frac{\rho_{\text{DM}}}{\rho_{\text{tot}}} h^2 \sim \frac{0.1 \text{ pb}}{\langle \sigma v \rangle}$$


$$\sim 0.1 \left( \frac{m_{\text{DM}}}{100 \text{ GeV}} \right)^2$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects



# 5. Prospects

## The many virtues of the WIMP



$$\Omega_{\text{DM}} h^2 \equiv \frac{\rho_{\text{DM}}}{\rho_{\text{tot}}} h^2 \sim \frac{0.1 \text{ pb}}{\langle\sigma v\rangle}$$


$$\sim 0.1 \left( \frac{m_{\text{DM}}}{100 \text{ GeV}} \right)^2$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Where are the WIMPs?

Direct detection







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects



# 5. Prospects

## Where are the WIMPs?

DM annihilation in the Galactic Center




K. Abazajian et al., PRD 102 (2020), 043012 (Fermi-LAT)




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Where are the WIMPs?

DM annihilation to neutrinos




C. Argüelles et al., arXiv:1912.09486 [hep-ph]




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs







# 4. Compact objects



# 5. Prospects

## Other evidence for Dark Matter?







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Beyond the WIMP




↓  
Feeble interactions =  
dependence on initial conditions




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

# Beyond the WIMP







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects



# 5. Prospects

## Inflating spacetime


Simplest incarnation: a slowly rolling scalar field in FRW spacetime,  $ds^2 = dt^2 - a(t)^2 dx^2$

$$\ddot{\Phi} + 3H\dot{\Phi} + V'(\Phi) = 0$$

$$H \equiv \frac{\dot{a}}{a} = \left( \frac{\rho_{\Phi}}{3M_P^2} \right)^{1/2} \quad \text{with}$$

$$\rho_{\Phi} = \frac{1}{2}\dot{\Phi}^2 + V(\Phi)$$


$$P_{\Phi} = \frac{1}{2}\dot{\Phi}^2 - V(\Phi)$$



When inflation ends, reheating begins




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects



# 5. Prospects

## The horizon problem


$$\Delta T/T \sim 10^{-5} \text{ in the CMB}$$



Credit: Héctor Ramírez




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects




# 5. Prospects

## The primordial fluctuations



Quantum fluctuations in  $\Phi$ ,  $g$ , are stretched by expansion

Planck TT, TE, EE + lowE + lensing + BK15 (■ 1σ, ■ 2σ)




Y. Akrami et al. [Planck], *Astron. Astrophys.* 641 (2020) A10

$$\mathcal{P}_{\mathcal{R}} = \frac{H_*^4}{4\pi^2 \dot{\Phi}_*^2} \left( \frac{k}{aH} \right)^{n_s - 1}$$

$$\mathcal{P}_{\mathcal{T}} = \frac{2H_*^2}{\pi^2} \left( \frac{k}{aH} \right)^{n_T}$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects




# 5. Prospects

## The primordial fluctuations



Quantum fluctuations in  $\Phi, g$ , are stretched by expansion



$$\mathcal{P}_{\mathcal{R}} = \frac{H_*^4}{4\pi^2 \dot{\Phi}_*^2} \left( \frac{k}{aH} \right)^{n_s - 1}$$


$$\mathcal{P}_{\mathcal{T}} = \frac{2H_*^2}{\pi^2} \left( \frac{k}{aH} \right)^{n_T}$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects



# 5. Prospects


## The vacuum can be excited!

$$A_\mu = (0, 0, 0, Et)$$


$$\omega^2(t) = (p_z + gEt)^2 + \mathbf{p}_T^2 + m^2$$

mixing of +/- frequency modes

particle production! (Schwinger effect)



During reheating, the inflaton provides an oscillating background



$$\mathcal{L}_\psi = y \Phi \bar{\psi} \psi \equiv m_\psi(t) \bar{\psi} \psi$$


$$\mathcal{L}_\chi = \frac{1}{2} \sigma \Phi^2 \chi^2 \equiv \frac{1}{2} m_\chi^2(t) \chi^2$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects






# 5. Prospects

## Scalar (p)reheating

$$\ddot{\chi}_p + 3H\dot{\chi}_p + \left[ \frac{p^2}{a^2} + m_\chi^2(t) \right] \chi_p = 0, \quad m_\chi^2(t) = \sigma\Phi^2 + m_{\chi,0}^2$$

Neglecting expansion,







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects



# 5. Prospects




## Scalar (p)reheating

$$\ddot{\chi}_p + 3H\dot{\chi}_p + \left[ \frac{p^2}{a^2} + m_\chi^2(t) \right] \chi_p = 0, \quad m_\chi^2(t) = \sigma\Phi^2 + m_{\chi,0}^2$$

With expansion,

$$X = a\chi$$

$$n_p = \frac{1}{2\omega_p} |\omega_p \chi_p - i\chi'|^2$$







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects






# 5. Prospects

## Fermion (p)reheating

$$\left[ i\gamma^\mu \partial_\mu + i\frac{3a'}{2a}\gamma^0 - am_\psi(\tau) \right] \psi = 0, \quad m_\psi^2(\tau) = (y\Phi + m_{\psi,0})^2$$

Neglecting expansion,\*







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects



# 5. Prospects

## Fermion (p)reheating

$$\left[ i\gamma^\mu \partial_\mu + i\frac{3a'}{2a}\gamma^0 - am_\psi(\tau) \right] \psi = 0, \quad m_\psi^2(\tau) = (y\Phi + m_{\psi,0})^2$$

Neglecting expansion,\*

\* $dt/d\tau = a$  and

$$\psi(\tau, \mathbf{x}) = a^{-3/2} \sum_{r=\pm} \int \frac{d^3 \mathbf{p}}{(2\pi)^{3/2}} e^{-i\mathbf{p}\cdot\mathbf{x}} \left[ u_p^{(r)}(\tau) \hat{a}_{\mathbf{p}}^{(r)} + v_p^{(r)}(\tau) \hat{b}_{-\mathbf{p}}^{(r)\dagger} \right]$$

with


$$u_p^{(r)}(\tau) = \begin{pmatrix} U_1(\tau) \xi_r(\mathbf{p}) \\ U_2(\tau) \frac{\boldsymbol{\sigma}\cdot\mathbf{p}}{p} \xi_r(\mathbf{p}) \end{pmatrix}$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects






# 5. Prospects

## Fermion (p)reheating

$$\left[ i\gamma^\mu \partial_\mu + i\frac{3a'}{2a}\gamma^0 - am_\psi(\tau) \right] \psi = 0, \quad m_\psi^2(\tau) = (y\Phi + m_{\psi,0})^2$$

With expansion,







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects




# 5. Prospects

## The perturbative (dissipative) picture

Reheating as the exchange of energy between two ideal fluids

$$T^{\mu\nu} = T_{\Phi}^{\mu\nu} + T_R^{\mu\nu} = \begin{pmatrix} \rho_{\Phi} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + \frac{1}{3} \begin{pmatrix} 3\rho_R & 0 & 0 & 0 \\ 0 & \rho_R & 0 & 0 \\ 0 & 0 & \rho_R & 0 \\ 0 & 0 & 0 & \rho_R \end{pmatrix}$$



$$\langle p_{\Phi} \rangle = \frac{1}{2} \langle \dot{\Phi}^2 + m_{\Phi}^2 \Phi^2 \rangle \simeq 0$$


(matter)




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects



# 5. Prospects


## The perturbative (dissipative) picture

Reheating as the exchange of energy between two ideal fluids


$$T^{\mu\nu} = T_{\Phi}^{\mu\nu} + T_R^{\mu\nu} = \begin{pmatrix} \rho_{\Phi} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + \frac{1}{3} \begin{pmatrix} 3\rho_R & 0 & 0 & 0 \\ 0 & \rho_R & 0 & 0 \\ 0 & 0 & \rho_R & 0 \\ 0 & 0 & 0 & \rho_R \end{pmatrix}$$

Conservation  $\nabla_{\mu} T^{\mu\nu} = 0,$

$$\dot{\rho}_R + 4H\rho_R = -(\dot{\rho}_{\Phi} + 3H\rho_{\Phi}) \equiv \Gamma_{\Phi}\rho_{\Phi}$$



$$\Gamma_{\Phi} = \frac{y^2}{8\pi} m_{\Phi}$$







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects




# 5. Prospects

## The perturbative (dissipative) picture


Reheating as the exchange of energy between two ideal fluids

$$T^{\mu\nu} = T_{\Phi}^{\mu\nu} + T_R^{\mu\nu} = \begin{pmatrix} \rho_{\Phi} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} + \frac{1}{3} \begin{pmatrix} 3\rho_R & 0 & 0 & 0 \\ 0 & \rho_R & 0 & 0 \\ 0 & 0 & \rho_R & 0 \\ 0 & 0 & 0 & \rho_R \end{pmatrix}$$







# 1. Beyond WIMPs



# 2. Inflation & reheating







# 3. FIMPs



# 4. Compact objects




# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating





# 3. FIMPs



# 4. Compact objects



# 5. Prospects



Early times:

field picture

(limited dissipation)

interactions  
(thermalization)

Late times:

fluid picture


(no fluctuations)




# 1. Beyond WIMPs



# 2. Inflation & reheating





# 3. FIMPs



# 4. Compact objects



# 5. Prospects



Early times:

field picture

(limited dissipation)

Schwinger-Keldysh / Kadanoff-Baym

Boltzmann

Late times:

fluid picture


(no fluctuations)




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## The phase space distribution

$$n_\psi = \int \frac{d^3\mathbf{p}}{(2\pi)^3} f_\psi(p, t)$$




$$\begin{aligned} \frac{\partial f_\psi}{\partial t} - H|\mathbf{p}| \frac{\partial f_\psi}{\partial |\mathbf{p}|} &= \frac{1}{2p_0} \int d\Pi |\mathcal{M}|_{\Phi \rightarrow \bar{\psi}\psi}^2 n_\Phi \delta^{(3)}(\mathbf{P}) + \mathcal{C}^{\text{int}}[f_\psi] \end{aligned}$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects



# 5. Prospects

## The phase space distribution

$$n_\psi = \int \frac{d^3\mathbf{p}}{(2\pi)^3} f_\psi(p, t)$$







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## The phase space distribution

$$n_\psi = \int \frac{d^3 \mathbf{p}}{(2\pi)^3} f_\psi(p, t)$$




$$C^{\text{int}}[f_\psi] = \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right|^2 + \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right|^2 + \dots$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects



# 5. Prospects

## The phase space distribution

$$n_\psi = \int \frac{d^3 \mathbf{p}}{(2\pi)^3} f_\psi(p, t)$$







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects



# 5. Prospects


Is a spin- $\frac{3}{2}$  dark matter particle the missing piece in the puzzle?

SM:

spin 0




spin 1/2



spin 1



spin 3/2



spin 2







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

Is a spin- $\frac{3}{2}$  dark matter particle the missing piece in the puzzle?




$$\mathcal{L} = -\frac{1}{2} \bar{\Psi}_\mu (i\gamma^{[\mu} \gamma^\nu \gamma^{\rho]} \partial_\rho + m_{3/2} \gamma^{[\mu} \gamma^{\nu]}) \Psi_\nu \quad (\text{Rarita-Schwinger})$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

Is a spin- $\frac{3}{2}$  dark matter particle the missing piece in the puzzle?



$$\mathcal{L} = -\frac{1}{2} \bar{\Psi}_\mu (i\gamma^{[\mu} \gamma^\nu \gamma^{\rho]} \partial_\rho + m_{3/2} \gamma^{[\mu} \gamma^{\nu]}) \Psi_\nu$$

$$+ y H \bar{\nu}_L \nu_R + \frac{M_R}{2} \bar{\nu}_R^c \nu_R \quad (\nu \text{ masses through see-saw})$$


$$+ y_\nu \Phi \bar{\nu}_R \nu_R \quad (\text{reheating})$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

Is a spin- $\frac{3}{2}$  dark matter particle the missing piece in the puzzle?



$$\mathcal{L} = -\frac{1}{2} \bar{\Psi}_\mu (i\gamma^{[\mu} \gamma^\nu \gamma^{\rho]} \partial_\rho + m_{3/2} \gamma^{[\mu} \gamma^{\nu]}) \Psi_\nu$$

$$+ y H \bar{\nu}_L \nu_R + \frac{M_R}{2} \bar{\nu}_R^c \nu_R$$

$$+ y_\nu \Phi \bar{\nu}_R \nu_R$$


$$+ i \frac{\alpha_1}{2M_P} \bar{\nu}_R \gamma^\mu [\gamma^\rho, \gamma^\sigma] \Psi_\mu F_{\rho\sigma}$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

Is a spin- $\frac{3}{2}$  dark matter particle the missing piece in the puzzle?




$$\begin{aligned}
 \mathcal{L} = & -\frac{1}{2} \bar{\Psi}_\mu (i\gamma^{[\mu} \gamma^\nu \gamma^{\rho]} \partial_\rho + m_{3/2} \gamma^{[\mu} \gamma^{\nu]}) \Psi_\nu \\
 & + y H \bar{\nu}_L \nu_R + \frac{M_R}{2} \bar{\nu}_R^c \nu_R \\
 & + y_\nu \Phi \bar{\nu}_R \nu_R \\
 & + i \frac{\alpha_1}{2M_P} \bar{\nu}_R \gamma^\mu [\gamma^\rho, \gamma^\sigma] \Psi_\mu F_{\rho\sigma} + i \frac{\alpha_2}{2M_P} i\sigma_2 (D^\mu H)^* \bar{L} \Psi_\mu + \dots
 \end{aligned}$$




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Scatterings and decays

$$\mathcal{L}_{3/2} = i \frac{\alpha_1}{2M_P} \bar{\nu}_R \gamma^\mu [\gamma^\rho, \gamma^\sigma] \Psi_\mu F_{\rho\sigma} + i \frac{\alpha_2}{2M_P} i\sigma_2 (D^\mu H)^* \bar{L} \Psi_\mu + \text{h.c.}$$




$$\Rightarrow \sigma(s) \propto \frac{\alpha_1^2 y^2 s^2}{m_{3/2}^2 M_R^2 M_P^2}$$


- Production peaked at high energies  $\rightarrow$  reheating
- $\Psi$  is never in thermal equilibrium (freeze-in)




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects




# 5. Prospects

## Scatterings and decays




$$\Omega_{3/2} \propto \alpha_1 \frac{m_\nu m_\Phi^2 T_{\text{reh}}^3}{m_{3/2} M_R t_{\text{therm}}}$$







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects




# 5. Prospects

## Scatterings and decays




$$\Omega_{3/2} \propto \alpha_1 \frac{m_\nu T_{\text{reh}}^5}{m_{3/2} M_R}$$







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects




# 5. Prospects

## Scatterings and decays




## Scattering






# How warm is out-of-equilibrium dark matter?


1. Beyond WIMPs



2. Inflation & reheating




3. FIMPs




4. Compact objects



5. Prospects



CDM




WDM (0.5 keV)




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects




# 5. Prospects

## How warm is out-of-equilibrium dark matter?







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects



# 5. Prospects

## How warm is out-of-equilibrium dark matter?

R. Murgia, V. Iršič and M. Viel, PRD 98 (2018), 083540




G. Ballesteros, MG and M. Pierre, JCAP 03 (2021), 101




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects




# 5. Prospects

Constraints:  $\Omega_{\text{DM}} + \gamma + \nu + \text{Lyman-}\alpha$



Scattering







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects





# 5. Prospects




## Production (via inflaton decay)

Via  $\alpha_1$ ,

$M_R \ll m_\Phi$ :



$M_R \gg m_\Phi$ :




(via  $\alpha_2$  are 2-loop suppressed)




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects




# 5. Prospects

Constraints:  $\Omega_{\text{DM}} + \gamma + \nu + \text{Lyman-}\alpha$




Inflaton decay







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects


## Beyond perturbation theory

Super-heavy dark matter (WIMPzillas, ...)



(but not thermally though)

E. Kolb, D. Chung and A. Riotto, AIP Conf. Proc. 484 (1999), 91




MG, K. Kaneta, Y. Mambrini and K. Olive, JCAP 04 (2021), 012




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Large metric fluctuations?



Credit: Naoya Kitajima

Metric preheating → overdensities → light PBHs → (Hawking) reheating

$M_{\text{BH}}$	$\tau_{\text{BH}}$
A man	$10^{-12}$ s
$10^{15}$ g	$10^{10}$ y
Earth	$10^{49}$ y
Sun	$10^{66}$ y
Milky Way	$10^{99}$ y


$$\frac{\delta\rho_C}{\rho} \gtrsim 0.5$$

$$M_{\text{BH}} \propto \rho_{\text{form}} / H_{\text{form}}^3 < 10^9 \text{ g}$$


$$\tau_{\text{BH}} \approx 10^{64} \left( \frac{M}{M_{\odot}} \right)^3 \text{ y}$$




# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects




# 5. Prospects

## Large metric fluctuations?




Credit: Naoya Kitajima




K. Jedamzik, M. Lemoine and J. Martin, JCAP 09 (2010), 034; **A. Gómez**, MG, G. Ballesteros, M. Pierre, *on it*




# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects





# 5. Prospects

## Fragmenting the condensate



“opening up” = attractive interaction


$$V = \frac{1}{2} m_\Phi^2 \Phi^2 + V_{NL}$$




Complex, U(1):	Q-Balls		Boson stars
Real:	Oscillons		Oscillatons




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Oscillaton collisions







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Oscillaton collisions







# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects


## Oscillaton collisions




LIGO range  $10^{-12}$  eV  $\lesssim m \lesssim 10^{-10}$  eV




# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects



# 5. Prospects

## Dark matter from inflation

ON THE CONCENTRATION OF RELIC MAGNETIC MONOPOLES IN THE UNIVERSE


Ya.B. ZELDOVICH and M.Yu. KHLOPOV

*Institute of Applied Mathematics, Academy of Sciences of the USSR, Moscow 125047, USSR*

PLB 79 (1978), 239

Inflation efficiently dilutes relics, dangerous or not


But inflation can lead to overdensities




G. Ballesteros and M. Taoso, PRD 97 (2018), 023501




# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects



# 5. Prospects

## Dark matter from inflation

### ON THE CONCENTRATION OF RELIC MAGNETIC MONOPOLES IN THE UNIVERSE


Ya.B. ZELDOVICH and M.Yu. KHLOPOV

*Institute of Applied Mathematics, Academy of Sciences of the USSR, Moscow 125047, USSR*

PLB 79 (1978), 239

Inflation efficiently dilutes relics, dangerous or not

But inflation can lead to overdensities




[github.com/bradkav/PBHbounds](https://github.com/bradkav/PBHbounds)




# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs







# 4. Compact objects



# 5. Prospects

## Warm inflation?



A. Berera, PRL 75 (1995), 3218

A. Berera, I. G. Moss and R. O. Ramos, Rept. Prog. Phys. 72 (2009), 026901

Inflaton slows down by thermal friction

$$\ddot{\Phi} + (3H + \Gamma)\dot{\Phi} + V_{\Phi} = 0$$

$$\dot{\rho}_r + 4H\rho_r - \Gamma\dot{\Phi}^2 = 0$$

Thermal noise is an extra source of density fluctuations

$$\delta\ddot{\Phi}_k + (3H + \Gamma)\delta\dot{\Phi}_k = \xi_k + \dots$$



Daniel Green @nu\_phases · 1 jun. ...

3/8 The inflaton needs to continually produce particles

Problem: most couplings that do this will destroy inflation / scale invariance

Worse problem: If you write a coupling that preserves scale invariance, it was shown you don't get normal friction


[arxiv.org/abs/1109.4192](https://arxiv.org/abs/1109.4192)




# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs



# 4. Compact objects




# 5. Prospects

## Warm inflation?


A way around? Warm Little Inflation ( $\Phi$  as a pseudo-Nambu-Goldstone boson)

Leftover  $\Phi$  as DM



J. Rosa and L. Ventura, PRL 122 (2019), 161301

PBH from WLI




M. Bastero-Gil and M. Díaz-Blanco, arXiv:2105.08045 [hep-ph]


A. Pérez, MG, G. Ballesteros, M. Pierre, J. Rey, on it

# Complexity in the early Universe


1. Beyond WIMPs



2. Inflation & reheating



3. FIMPs




4. Compact objects



5. Prospects


## Particle theory

- SM UV completions  $N_F \gg 1$
- Coupling to  $\Phi$  weakly constrained
- Non-trivial field manifolds







# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## Complexity in the early Universe




R. Flauger et al., JCAP 10 (2017), 058  
 J. Ellis, MG et al., JCAP 07 (2017), 006


$$m_{\chi}^2(t) = m_{\chi,0}^2 + g^2 \sum_i (\Phi(t) - \Phi_i)^2 + \dots$$

R. Easter and L. McAllister, JCAP 05 (2006), 018  
 D. Green et al., PRD 80 (2009), 063533  
 Y. Nakai, R. Namba and Z. Wang, JHEP 12 (2020), 170


# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs





# 4. Compact objects



# 5. Prospects

## Complexity in the early Universe




$$m_{\chi}^2(t) = m_{\chi,0}^2 + g^2 \sum_i (\Phi(t) - \Phi_i)^2 + \dots$$


$$(\zeta \simeq \mathcal{R})$$




# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects




# 5. Prospects

## A (conformal) spectator in dS


MG, M. Amin, S. Carlsten and D. Green, JCAP 05 (2019), 012




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects

## A (conformal) spectator in dS







# 1. Beyond WIMPs




# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects




# 5. Prospects

## A stochastically sourced power spectrum


MG, M. A. Amin and D. Green, JCAP 06 (2020), 039




# 1. Beyond WIMPs



# 2. Inflation & reheating




# 3. FIMPs



# 4. Compact objects




# 5. Prospects







# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs






# 4. Compact objects




# 5. Prospects


## Prospects




# 1. Beyond WIMPs



# 2. Inflation & reheating




# 3. FIMPs




# 4. Compact objects




# 5. Prospects




O. Newton et al., arXiv:2011.08865



CTA, JCAP 01 (2021), 057




gwplotter.com




Simons Observatory, JCAP 02 (2019), 056




# 1. Beyond WIMPs



# 2. Inflation & reheating



# 3. FIMPs




# 4. Compact objects



# 5. Prospects


## Reheating + BSM is not always simple

Multifield effects




J. Ellis, MG, N. Nagata, D. Nanopoulos and K. Olive, JCAP 07 (2017), 006

In-medium effects



V. Rychkov and A. Strumia, PRD 75 (2007), 075011


More general potentials




MG, K. Kaneta, Y. Mambrini and K. A. Olive, PRD 101 (2020), 123507 ;

MG, K. Kaneta, Y. Mambrini and K. A. Olive, JCAP 04 (2021), 012


## 1. Beyond WIMPs



## 2. Inflation & reheating



## 3. FIMPs




## 4. Compact objects



## 5. Prospects

## Conclusion

- The light beyond the WIMP: the Big Bang itself?
- Dark matter as a probe of early dynamics
- Indirect connection with low energy physics (neutrinos, CMB, proton decay, ...)
- Same ideas, different applications (condensed matter, baryon asymmetry, finance, ...)



谢谢!